

Results from POLARBEAR and plans for the Simons Array

Nils Halverson
University of Colorado – Boulder
for the POLARBEAR Collaboration



POLARBEAR Collaboration

UC Berkeley

Brian Barch
Ari Cukierman
Tijmen de Haan
Josquin Errard
Neil Goeckner-Wald
Charles Hill
William Holzapfel
Yasuto Hori
Oliver Jeong
Adrian Lee
Mike Myers
Chris Raum
Paul Richards
Blake Sherwin
Ian Shirley
Bryan Steinbach
Nathan Whitehorn
Oliver Zahn



Lawrence Berkeley NL

Julian Borrill
Reijo Keskitalo
Theodore Kisner
Akito Kusaka
Eric Linder
Aritoki Suzuki



UC San Diego

Chris Aleman
Kam Arnold
Matt Atlas
Darcy Barron
Tucker Elleflot
George Fuller
Logan Howe
Jon Kaufman
Kavon Kazemzadeh
Brian Keating
David Leon
Lindsay Lowry
Frederick Matsuda
Martin Navaroli
Hans Paar
Gabriel Rebeiz
Praween Siritanasak
Nathan Stebor
Brandon Wilson
Amit Yadav



CU Boulder

Nils Halverson
Greg Jaehnig
David Schenck



KEK

Yuji Chinone
Masaya Hasegawa
Kaori Hattori
Masashi Hazumi
Takahiro Okamura
Jun-ichi Suzuki
Osamu Tajima
Satoru Takakura
Takayuki Tomaru



SOKENDAI

Yoshiki Akiba
Yuki Inoue
Yuuko Segawa



Laboratoire Astroparticule & Cosmologie

Maude Le Jeune
Julien Peloton
Davide Poletti
Radek Stomp



Argonne NL

Amy Bender



Cardiff U.

Peter Ade



McGill University

Matt Dobbs
Adam Gilbert
Josh Montgomery
Graeme Smecher



Dalhousie

Scott Chapman
Colin Ross
Kaja Rotermund
Alexei Tikhomirov



Kavli IPMU

Fumiya Irie
Nobuhiko Katayama
Kuniyoshi Mizukami
Haruki Nishino



Imperial College

Anne Ducout
Stephen Feeney
Andrew Jaffe



U. Melbourne

Christian Reichardt



SISSA

Carlo Baccigalupi
Giulio Fabbian
Giuseppe Puglisi



JAXA

Tomotake Matsumura



UC Irvine

Chang Feng



National Institute for Fusion Science

Suguru Takada



Cardiff University

Peter Ade



NASA Goddard

Nathan Miller



Princeton

Zigmund Kermish



Católica (PUC)


David Boettger
Rolando Dunner





And many more in years past




Outline



- Scientific goals
- The POLARBEAR instrument
- First season results
- The next generation instrument: PB2/Simons Array





Predicted Polarization Power Spectrum



Lensing potential map from correlating across angular scales in T,E,B

- Reconstruct gravitational potential
- Lensing potential power spectrum:
 - Neutrino mass
 - Dark energy
- Deflection map:
 - De-lensing




POLARBEAR Instrument Design: Telescope

Huan Tran Telescope

Angular
resolution to
characterize
lensing,
3.5' @ 150
GHz

Off-axis
Gregorian-
Dragone:
-Low cross
polarization
-Large field of
view
-Large
throughput




Elevation: 5200 m

Shielding to
reduce sidelobe
response


Cryogenic
Receiver
enclosure

POLARBEAR Cryogenic Receiver




POLARBEAR: Control of systematic instrumental effects

- Instantaneous measurement of polarization with each pixel
- Modulation of CMB Polarization:
 - Half-wave plate
 - Apparent sky rotation
- Low sidelobe response
 - Co-moving baffling
 - Scan strategy allows ground removal



Instrumental Bias Estimation

End-to-end simulations using measured instrument characteristics and cross-checks




Confirmed all known systematics are much smaller than statistical uncertainty before “unblinding” the spectrum

POLARBEAR First Season Observations


Stokes Q map

μK_{CMB}



Stokes U map

μK_{CMB}




- Three fields, 24.5 deg^2 total sky area
- Map depth: 5.5, 8.5, 7.6 $\mu\text{K-arcmin}$
- Array sensitivity: $23 \mu\text{K s}^{-1/2}$
- 13 calendar months of observations (May 2012 - June 2013)

First-Season POLARBEAR Results: Three B-mode Analyses


Galaxy cross-correlation:

PRL 112, 131302 (2014)




Deflection power spectrum:

PRL 113, 021301 (2014)



Angular power spectrum: C_{ℓ}^{BB}

ApJ 794, 171 (2014)




All Spectra available on NASA LAMBDA site

All analysis use 13 calendar months of observations (May 2012 - June 2013)

B-modes: Galaxy cross correlation


polarization lensing \times CIB



- 4.0σ evidence for gravitational lensing of CMB polarization
- Consistent with SPTpol measurement

B-modes: deflection power spectrum

- 4.2 sigma rejection of “no lensing B -modes”
- First reported CMB-polarization-only C_L^{dd} spectrum
- Polarization will be the most sensitive measure of C_L^{dd}
 - Groundwork for neutrino masses, dark energy



Green: $\langle EBEB \rangle$

Blue: $\langle EEEB \rangle$


Red: Combined

B-modes: Power Spectrum C_{ℓ}^{BB}

97.2% rejection of “no lensing B -modes”


- including possible systematic instrumental biases
- Including estimate of foreground bias

Combining three published results: 4.7σ rejection of “no lensing B -modes”



Amplitude of lensing compared to Λ CDM:
 1.12 ± 0.61 (*stat*) $^{+0.04}_{-0.12}$ (*sys*) ± 0.07 (*mult*)



B-modes: Power Spectrum C_ℓ^{BB}





POLARBEAR -> Simons Array

- Simons Array
 - Sensitivity: 22,768 total detectors
 - Resolution: 4' sys. error advantage, measure lensing
 - Multi-chroic pixels with 90/150 and 150/220 GHz frequency coverage
 - Wide sky survey ($f_{\text{sky}} = 65\%$)
 - Scalability: Future expansion possible (more telescopes and frequencies)

POLARBEAR-2: Sinuous Focal Plane






- POLARBEAR-1
- 1,274 Bolometers
- 150 GHz
- 22 cm diameter
- 4" wafers



- POLARBEAR-2
- 7,588 Bolometers
- 95/150 GHz pixels
- 36.5 cm diameter
- 6" wafers

POLARBEAR-2 Receiver

365 mm diameter focal plane
7,588 bolometers
2-band Pixel



Lab tests underway of:

- Assembled cryogenic receiver
- Prototype detector sub-arrays
- 40x frequency multiplexed readout

The Simons Array

Leverage POLARBEAR experience to rapidly increase sensitivity





2014-2015: Construct two more telescopes

2015: POLARBEAR-2 (95 GHz / 150 GHz) deploys onto first new telescope


2016: A copy of POLARBEAR-2 deploys onto second new telescope

2016: A 150 GHz / 220 GHz receiver replaces POLARBEAR-1 on the original telescope

3 receivers (22,764 bolometers) observing at 95,150,220 GHz
All hardware funded by the Simons Foundation, MEXT, and NSF



Simons Array Sensitivity




Foreground rejection
with 95/150/220 GHz,
Planck, & C-BASS

- $\sigma(r=0.1) = 6 \times 10^{-3}$
- $\sigma(n_s) = 6 \times 10^{-3}$

Neutrino Mass:

- $\sigma(\Sigma m_\nu) = 40 \text{ meV}$
(w/DESI BAO)

In Summary...



- POLARBEAR-1: Three analyses showing B-modes (CMB alone: 4.7σ rejection of no lensing B-modes)
 - CIB cross-correlation
 - Deflection power spectrum
 - C_l^{BB}
- Future: multi-chroic arrays in the Simons Array
 - Increased sensitivity
 - Foreground mitigation

